

POSHERSTNIK, M.Yu., kand. tekhn. nauk; SALYUTINA, M.A., inzh.

Using powered germanium rectifiers in marine electric systems.
Sudostroenie 24 no.10:33-35 0 '58. (MIRA 11:12)
(Electricity on ships)

POSHERSTNIK, M.Yu., kand.tekhn.nauk

Using cables with mineral insulation in shipbuilding. Snistrenis
24 no.5:58-59 My '58. (MIRA 11:6)
(Electric cables) (Electricity on ships)

PUSHKAR', L.N.; POSHEVAYA, V.P.; GERASIMOVA, L.I.; TROITSKIY, V.B.

Clinico-experimental study of the hydrolyzate aminophaseol.
Vest. khir. 70 no.6:26-29 Ja'63 (MIRA 16:12)

1. Iz Tsentral'nogo ordena Lenina instituta gematologii i
perelivaniya krovi (dir. - prof. A.A. Bagdasarov [deceased]).
Adres avtorov: Moskva, Novo-Zykovskiy proyезд, 4, Tsentral'-
nyy institut gematologii i perelivaniya krovi.

POSHITNOV, V.N.

5

AGAFANOV, A.V., RYSAKOV, M.V., COLUSHESYN, D.L., GUSELKOVA, YE.A.,
ALFIMOVA, YE.A., POSHITNOV, V.N.,

Gewinnung von Motorölen aus schwefelhaltigen Rohölen durch
Hydrierung.

Report to be submitted for the Symposium Lubricants and
Lubrication, Dresden, 27-30 June 1961

POSHKA, A.L., Cand Tech Sci - -(diss) "Determination of the maximum
expenditure of water for the calculation of water intakes and
offtakes in the Latvian SSR," Kaunas, 1960, 20 pp (Kaunas Polytechnical
Institute) (KL, 36-60, 115)

POSHKA, A.I.; FEDORCHUK, V.P., kand. geologo-mineral. nauk

Ancient mercury workings in southern Fergana. Priroda 48 no.5:
108-109 My '59. (MIRA 12:5)

1.Omskiy oblastnoy krayevedcheskiy muzey (for Poshka). 2.Khaydarkanskaya
geologorazvedochnaya partiya tresta "Sredaztsvetmetrazvedka", Omskaya
oblasti Kirgiskoy SSR (for Federchuk).
(Fergana--Mercury mines and mining)

18(5)

SOV/26-59-5-29/47

AUTHORS: Poshka, A.I. and Fedorchuk, V.P., Candidate of Geomineralogical Sciences

TITLE: Ancient Mercury Mines in South Fergana

PERIODICAL: Priroda, 1959, Nr 5, pp 108 - 109 (USSR)

ABSTRACT: The authors refer to the Tadzhik-Pamir 1926-37 expedition of the Academy of Sciences of the USSR, when D.I. Shcherbakov defined the mercury antimony belt in that area, known from ancient times. The remains of the old mining sites (subterranean passages and halls) were found and proofs obtained that considerable quantities of cinnabar and mercury were exported from this province to India and China.

Card 1/2

MATSEVICIUS, I.I.; POSHKA, A.L.

Determination of the reduction coefficient in empiric formulas for
calculating maximum discharges. Meteor. i gidrol. no.3:43-44 Mr
'56. (Stream measurements) (MIRA 9:7)

KUGATOVA-SHENYAKINA, G.P.; POSHKENE, R.A.

Synthesis of Δ^6 -cyclohexene-aldehydes from isomeric Δ^3 -aldehydes.
Zhur.ob.khim. 32 no.8:2461-2464, Ag '62. (MIRA 15:9)
(Benzaldehyde)

KUGATOVA-SHEMYAKINA, G.P.; LAUMYANSKAS, G.A.; KRASIL'NIKOVA, G.K.;
MOZOLIS, V.V.; KAYKARIS, P.A.; POSEKENE, R.A.

Ethynylation of ionone analogs. Zhur.ob.khim. 32 no.8:2455-2461
Ag '62. (MIRA 15:9)

(Ionone) (Ethynylation)

KUGATOVA, G.P. [Kugatova, G.]; POSHKENE, R.A. [Poshkiene, R.]

Aromatization of 2-methyl-, 4-methyl- Δ^3 -tetrahydrobenzaldehydes.
Liet ak darbai B no.2:157-161 '60. (EEAI 10:1)

1. Institut khimii i khimicheskoy tekhnologii Akademii nauk
Litovskoy SSR
(Aromatization) (Dimethylcyclohexenealdehyde)

MINENKO, V.A.; PEYSHEV, G.P.; KURILOV, P.G.; VERZHIKOVSKAYA, L.G.;
VASIL'YEVA, S.M.; POSHKREBNEV, V.A.

Potentialities for increasing the output of open-hearth
furnace plants now in operation. Stal' 23 [i.e. 24] no.4:
309-313 Ap '84. (MIRA 17:8)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut organizatsii
proizvodstva i truda chernoy metallurgii.

MINENKO, V.A.; ALEKSANDROV, A.A.; SVETS, V.Ye.; BORZENKO, V.P.; KURILOV,
P.G.; KHAZANOVICH, N.L.; Prinimali uchastiye: POPOV, A.I.;
KONOVALOV, A.N.; TERTYCHNAYA, I.Yu.; POCHIZHENY, Y.P.;
DMITRIYEVA, S.M.; KORNILOVA, A.V.

Work organization in the section, of metal feed to blooming
mills. Met. i gornorud. prom. no.2:67-68 Mr-Ap '64.

(MIRA 17:9)

15

PROCESSES AND PROPERTIES INDEX

Colorimetric Determination of Copper With the Aid of Dimethylglyoxime. (In Russian.) V. M. Babkova, M. E. Leontin, and K. I. Litvin. Zhurnal Analiticheskoi Khimii (Journal of Analytical Chemistry), v. 3, May-June 1948, p. 161-166.

The methods of Goethals and of Clarke and Jones were studied. Certain modifications of the latter were developed, resulting in more stable coloration. The nature of the reaction was studied.

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

13000 13100 13200 13300 13400 13500 13600 13700 13800 13900 14000 14100 14200 14300 14400 14500 14600 14700 14800 14900 15000 15100 15200 15300 15400 15500 15600 15700 15800 15900 16000 16100 16200 16300 16400 16500 16600 16700 16800 16900 17000 17100 17200 17300 17400 17500 17600 17700 17800 17900 18000 18100 18200 18300 18400 18500 18600 18700 18800 18900 19000 19100 19200 19300 19400 19500 19600 19700 19800 19900 20000 20100 20200 20300 20400 20500 20600 20700 20800 20900 21000 21100 21200 21300 21400 21500 21600 21700 21800 21900 22000 22100 22200 22300 22400 22500 22600 22700 22800 22900 23000 23100 23200 23300 23400 23500 23600 23700 23800 23900 24000 24100 24200 24300 24400 24500 24600 24700 24800 24900 25000 25100 25200 25300 25400 25500 25600 25700 25800 25900 26000 26100 26200 26300 26400 26500 26600 26700 26800 26900 27000 27100 27200 27300 27400 27500 27600 27700 27800 27900 28000 28100 28200 28300 28400 28500 28600 28700 28800 28900 29000 29100 29200 29300 29400 29500 29600 29700 29800 29900 30000 30100 30200 30300 30400 30500 30600 30700 30800 30900 31000 31100 31200 31300 31400 31500 31600 31700 31800 31900 32000 32100 32200 32300 32400 32500 32600 32700 32800 32900 33000 33100 33200 33300 33400 33500 33600 33700 33800 33900 34000 34100 34200 34300 34400 34500 34600 34700 34800 34900 35000 35100 35200 35300 35400 35500 35600 35700 35800 35900 36000 36100 36200 36300 36400 36500 36600 36700 36800 36900 37000 37100 37200 37300 37400 37500 37600 37700 37800 37900 38000 38100 38200 38300 38400 38500 38600 38700 38800 38900 39000 39100 39200 39300 39400 39500 39600 39700 39800 39900 40000 40100 40200 40300 40400 40500 40600 40700 40800 40900 41000 41100 41200 41300 41400 41500 41600 41700 41800 41900 42000 42100 42200 42300 42400 42500 42600 42700 42800 42900 43000 43100 43200 43300 43400 43500 43600 43700 43800 43900 44000 44100 44200 44300 44400 44500 44600 44700 44800 44900 45000 45100 45200 45300 45400 45500 45600 45700 45800 45900 46000 46100 46200 46300 46400 46500 46600 46700 46800 46900 47000 47100 47200 47300 47400 47500 47600 47700 47800 47900 48000 48100 48200 48300 48400 48500 48600 48700 48800 48900 49000 49100 49200 49300 49400 49500 49600 49700 49800 49900 50000 50100 50200 50300 50400 50500 50600 50700 50800 50900 51000 51100 51200 51300 51400 51500 51600 51700 51800 51900 52000 52100 52200 52300 52400 52500 52600 52700 52800 52900 53000 53100 53200 53300 53400 53500 53600 53700 53800 53900 54000 54100 54200 54300 54400 54500 54600 54700 54800 54900 55000 55100 55200 55300 55400 55500 55600 55700 55800 55900 56000 56100 56200 56300 56400 56500 56600 56700 56800 56900 57000 57100 57200 57300 57400 57500 57600 57700 57800 57900 58000 58100 58200 58300 58400 58500 58600 58700 58800 58900 59000 59100 59200 59300 59400 59500 59600 59700 59800 59900 60000 60100 60200 60300 60400 60500 60600 60700 60800 60900 61000 61100 61200 61300 61400 61500 61600 61700 61800 61900 62000 62100 62200 62300 62400 62500 62600 62700 62800 62900 63000 63100 63200 63300 63400 63500 63600 63700 63800 63900 64000 64100 64200 64300 64400 64500 64600 64700 64800 64900 65000 65100 65200 65300 65400 65500 65600 65700 65800 65900 66000 66100 66200 66300 66400 66500 66600 66700 66800 66900 67000 67100 67200 67300 67400 67500 67600 67700 67800 67900 68000 68100 68200 68300 68400 68500 68600 68700 68800 68900 69000 69100 69200 69300 69400 69500 69600 69700 69800 69900 70000 70100 70200 70300 70400 70500 70600 70700 70800 70900 71000 71100 71200 71300 71400 71500 71600 71700 71800 71900 72000 72100 72200 72300 72400 72500 72600 72700 72800 72900 73000 73100 73200 73300 73400 73500 73600 73700 73800 73900 74000 74100 74200 74300 74400 74500 74600 74700 74800 74900 75000 75100 75200 75300 75400 75500 75600 75700 75800 75900 76000 76100 76200 76300 76400 76500 76600 76700 76800 76900 77000 77100 77200 77300 77400 77500 77600 77700 77800 77900 78000 78100 78200 78300 78400 78500 78600 78700 78800 78900 79000 79100 79200 79300 79400 79500 79600 79700 79800 79900 80000 80100 80200 80300 80400 80500 80600 80700 80800 80900 81000 81100 81200 81300 81400 81500 81600 81700 81800 81900 82000 82100 82200 82300 82400 82500 82600 82700 82800 82900 83000 83100 83200 83300 83400 83500 83600 83700 83800 83900 84000 84100 84200 84300 84400 84500 84600 84700 84800 84900 85000 85100 85200 85300 85400 85500 85600 85700 85800 85900 86000 86100 86200 86300 86400 86500 86600 86700 86800 86900 87000 87100 87200 87300 87400 87500 87600 87700 87800 87900 88000 88100 88200 88300 88400 88500 88600 88700 88800 88900 89000 89100 89200 89300 89400 89500 89600 89700 89800 89900 90000 90100 90200 90300 90400 90500 90600 90700 90800 90900 91000 91100 91200 91300 91400 91500 91600 91700 91800 91900 92000 92100 92200 92300 92400 92500 92600 92700 92800 92900 93000 93100 93200 93300 93400 93500 93600 93700 93800 93900 94000 94100 94200 94300 94400 94500 94600 94700 94800 94900 95000 95100 95200 95300 95400 95500 95600 95700 95800 95900 96000 96100 96200 96300 96400 96500 96600 96700 96800 96900 97000 97100 97200 97300 97400 97500 97600 97700 97800 97900 98000 98100 98200 98300 98400 98500 98600 98700 98800 98900 99000 99100 99200 99300 99400 99500 99600 99700 99800 99900 100000 100100 100200 100300 100400 100500 100600 100700 100800 100900 101000 101100 101200 101300 101400 101500 101600 101700 101800 101900 102000 102100 102200 102300 102400 102500 102600 102700 102800 102900 103000 103100 103200 103300 103400 103500 103600 103700 103800 103900 104000 104100 104200 104300 104400 104500 104600 104700 104800 104900 105000 105100 105200 105300 105400 105500 105600 105700 105800 105900 106000 106100 106200 106300 106400 106500 106600 106700 106800 106900 107000 107100 107200 107300 107400 107500 107600 107700 107800 107900 108000 108100 108200 108300 108400 108500 108600 108700 108800 108900 109000 109100 109200 109300 109400 109500 109600 109700 109800 109900 110000 110100 110200 110300 110400 110500 110600 110700 110800 110900 111000 111100 111200 111300 111400 111500 111600 111700 111800 111900 112000 112100 112200 112300 112400 112500 112600 112700 112800 112900 113000 113100 113200 113300 113400 113500 113600 113700 113800 113900 114000 114100 114200 114300 114400 114500 114600 114700 114800 114900 115000 115100 115200 115300 115400 115500 115600 115700 115800 115900 116000 116100 116200 116300 116400 116500 116600 116700 116800 116900 117000 117100 117200 117300 117400 117500 117600 117700 117800 117900 118000 118100 118200 118300 118400 118500 118600 118700 118800 118900 119000 119100 119200 119300 119400 119500 119600 119700 119800 119900 120000 120100 120200 120300 120400 120500 120600 120700 120800 120900 121000 121100 121200 121300 121400 121500 121600 121700 121800 121900 122000 122100 122200 122300 122400 122500 122600 122700 122800 122900 123000 123100 123200 123300 123400 123500 123600 123700 123800 123900 124000 124100 124200 124300 124400 124500 124600 124700 124800 124900 125000 125100 125200 125300 125400 125500 125600 125700 125800 125900 126000 126100 126200 126300 126400 126500 126600 126700 126800 126900 127000 127100 127200 127300 127400 127500 127600 127700 127800 127900 128000 128100 128200 128300 128400 128500 128600 128700 128800 128900 129000 129100 129200 129300 129400 129500 129600 129700 129800 129900 130000 130100 130200 130300 130400 130500 130600 130700 130800 130900 131000 131100 131200 131300 131400 131500 131600 131700 131800 131900 132000 132100 132200 132300 132400 132500 132600 132700 132800 132900 133000 133100 133200 133300 133400 133500 133600 133700 133800 133900 134000 134100 134200 134300 134400 134500 134600 134700 134800 134900 135000 135100 135200 135300 135400 135500 135600 135700 135800 135900 136000 136100 136200 136300 136400 136500 136600 136700 136800 136900 137000 137100 137200 137300 137400 137500 137600 137700 137800 137900 138000 138100 138200 138300 138400 138500 138600 138700 138800 138900 139000 139100 139200 139300 139400 139500 139600 139700 139800 139900 140000 140100 140200 140300 140400 140500 140600 140700 140800 140900 141000 141100 141200 141300 141400 141500 141600 141700 141800 141900 142000 142100 142200 142300 142400 142500 142600 142700 142800 142900 143000 143100 143200 143300 143400 143500 143600 143700 143800 143900 144000 144100 144200 144300 144400 144500 144600 144700 144800 144900 145000 145100 145200 145300 145400 145500 145600 145700 145800 145900 146000 146100 146200 146300 146400 146500 146600 146700 146800 146900 147000 147100 147200 147300 147400 147500 147600 147700 147800 147900 148000 148100 148200 148300 148400 148500 148600 148700 148800 148900 149000 149100 149200 149300 149400 149500 149600 149700 149800 149900 150000 150100 150200 150300 150400 150500 150600 150700 150800 150900 151000 151100 151200 151300 151400 151500 151600 151700 151800 151900 152000 152100 152200 152300 152400 152500 152600 152700 152800 152900 153000 153100 153200 153300 153400 153500 153600 153700 153800 153900 154000 154100 154200 154300 154400 154500 154600 154700 154800 154900 155000 155100 155200 155300 155400 155500 155600 155700 155800 155900 156000 156100 156200 156300 156400 156500 156600 156700 156800 156900 157000 157100 157200 157300 157400 157500 157600 157700 157800 157900 158000 158100 158200 158300 158400 158500 158600 158700 158800 158900 159000 159100 159200 159300 159400 159500 159600 159700 159800 159900 160000 160100 160200 160300 160400 160500 160600 160700 160800 160900 161000 161100 161200 161300 161400 161500 161600 161700 161800 161900 162000 162100 162200 162300 162400 162500 162600 162700 162800 162900 163000 163100 163200 163300 163400 163500 163600 163700 163800 163900 164000 164100 164200 164300 164400 164500 164600 164700 164800 164900 165000 165100 165200 165300 165400 165500 165600 165700 165800 165900 166000 166100 166200 166300 166400 166500 166600 166700 166800 166900 167000 167100 167200 167300 167400 167500 167600 167700 167800 167900 168000 168100 168200 168300 168400 168500 168600 168700 168800 168900 169000 169100 169200 169300 169400 169500 169600 169700 169800 169900 170000 170100 170200 170300 170400 170500 170600 170700 170800 170900 171000 171100 171200 171300 171400 171500 171600 171700 171800 171900 172000 172100 172200 172300 172400 172500 172600 172700 172800 172900 173000 173100 173200 173300 173400 173500 173600 173700 173800 173900 174000 174100 174200 174300 174400 174500 174600 174700 174800 174900 175000 175100 175200 175300 175400 175500 175600 175700 175800 175900 176000 176100 176200 176300 176400 176500 176600 176700 176800 176900 177000 177100 177200 177300 177400 177500 177600 177700 177800 177900 178000 178100 178200 178300 178400 178500 178600 178700 178800 178900 179000 179100 179200 179300 179400 179500 179600 179700 179800 179900 180000 180100 180200 180300 180400 180500 180600 180700 180800 180900 181000 181100 181200 181300 181400 181500 181600 181700 181800 181900 182000 182100 182200 182300 182400 182500 182600 182700 182800 182900 183000 183100 183200 183300 183400 183500 183600 183700 183800 183900 184000 184100 184200 184300 184400 184500 184600 184700 184800 184900 185000 185100 185200 185300 185400 185500 185600 185700 185800 185900 186000 186100 186200 186300 186400 186500 186600 186700 186800 186900 187000 187100 187200 187300 187400 187500 187600 187700 187800 187900 188000 188100 188200 188300 188400 188500 188600 188700 188800 188900 189000 189100 189200 189300 189400 189500 189600 189700 189800 189900 190000 190100 190200 190300 190400 190500 190600 190700 190800 190900 191000 191100 191200 191300 191400 191500 191600 191700 191800 191900 192000 192100 192200 192300 192400 192500 192600 192700 192800 192900 193000 193100 193200 193300 193400 193500 193600 193700 193800 193900 194000 194100 194200 194300 194400 194500 194600 194700 194800 194900 195000 195100 195200 195300 195400 195500 195600 195700 195800 195900 196000 196100 196200 196300 196400 196500 196600 196700 196800 196900 197000 197100 197200 197300 197400 1975

POSHKURLAT, A.P. (Moskva)

Analysis of the developmental rhythm of the cowslip primrose.
Bot.zhur. 47 no.2:262-267 F '62. (MIRA 15:3)
(Primroses)

POSHKURLAT, A.P.

Biology of the development of *Potentilla erecta* (L.) Ramps.
Trudy Len. khim.-farm. inst. 12:323-333 '61. (MIRA 15:3)

1. Kafedra botaniki Moskovskogo ordena Lenina meditsinskogo
instituta imeni I.M. Sechenova.

(CINQUEFOIL)

POSHKURLAT, A.P.

Protect medicinal plants. Apt. delo 11 no.623-25 K-D'62
(MIRA 1787)

1. Farmatsevticheskiy fakul'tet I Moskovskogo ordena Lenina
meditsinskogo instituta.

POSHKURLAT, A.P.

Changes occurring with age in the morphological features of *Potentilla erecta* (L.) Hampe [with summary in English]. BiolMOIP.Otd.
biol. 63 no.3:113-126 My-Je '58. (MIRA 12:3)
(GROWTH (PLANTS)) (CINQUEFOIL)

Country : USSR

Category: Cultivated Plants. Medicinal. Essential Oil
Bearing. Toxins.

M

Abs Jour: RZhBiol., No 22, 1958, No 100504

Author : Poshkurlat, A. P.

Inst : Moscow Pharmaceutical Inst.

Title : Anatomical Structure of the Cowslip Primrose
(*Primula veris* L.)

Orig Pub: Sb. nauchn. rabot. Mosk. farmatsevt. in-t, 1957,
1, 229-241

Abstract: Characteristic anatomical features of the
rhizome, adventitious root, the leaf, its
petiole, pedicel, peduncle and of the bract
are described. Changes are pointed out which

Card : 1/2

POSHKUS, D. P.

AUTHORS: Avgul', N. N., Isirikyan, A. A., 62-11-4/29
Kiselev, A. V., Lygina, I. A., Poshkus, D. P.

TITLE: Adsorption Equilibria and the Energy of Adsorption
Powers (Adsorbtsionnyye ravnovesiya i energiya
adsorbtsionnykh sil).

PERIODICAL: Izvestiya AN SSSR, Otdel. Khim. Nauk, 1957, Nr 11,
pp. 1314-1327 (USSR)

ABSTRACT: Here the theoretical and experimental investigation of the
adsorption powers in physical adsorption, mainly of complicated
non-polar molecules with adsorbents of an atomic and ionic
lattice, is brought. The results of the theoretical
computation are compared with the measurings of the dif-
ferential heats of the adsorption. Here a method for the
computation of the adsorption energy of non-polar molecules
with regard to three terms in the potential of the dispersion
powers with constants, which are computed by means of
polarizability and magnetization-coefficients, was worked
out. With it the induction potential by the average
polarizability of the adsorbed substance and the average
electrostatic field of the adsorbent was taken into
consideration. Furthermore the push-off potential with a

Card 1/3

Adsorption Equilibria and the Energy of Adsorption Powers 62-11-4/29

constant in the exponent, which is computed from the individual constants of the adsorbent, is taken into consideration. Finally all interactions of the given power center of the molecule of the adsorbed substance are added up with all adsorbent-lattice centers. The push-off constant before the exponential function is determined from the condition of the minimum of total energy of all interactions in an equilibrium distance from the adsorbent-surface. It is shown that the computed adsorption energy amounts of inert gases, nitrogen and 13 hydrocarbons of different structure (normal and isomeric alkanes, alcene, aromatical ones) on graphite are similar to the measured adsorption heats on graphited soot. It is shown that the computed adsorption energy amounts of the n-alkanes, of the benzene and toluene on magnesium oxide are also similar to the measured adsorption-heats. Furthermore it is shown, that in the case of an adsorption on graphite the amounts of the first, second and third term of the energy of dispersion powers and the absolute amount of the push-off energy were 90-95, or 5-10, or 0.5 - 1, or 35 - 40 % respectively of the total energy of dispersion powers in the investigated

Card 2/3

POSHKUS, D.P., Cand. Chem Sci—(disc) "Adsorption of vapors of benzene
and n. hexane on ^{chromium}oxide and hydroxide ~~of H₂~~." Mos, 1958. 19 pp (Acad of Sci
USSR. Inst of Physical Chemistry), 150 copies (ML,47-58,131)

-17-

POSHKUS, D. P.

62-58-4-29/32

AUTHORS: Kiselev, A. V., Poshekus, D. P.
 TITLE: Letter to the Editor (Pis'ma redaktoru)
 PERIODICAL: Izvestiya Akademii Nauk SSSR, Otdeleniye Khimicheskikh Nauk,
 1958, Nr 4, p . 520 (USSR)
 ABSTRACT: In an article by A.V. Kiselev, DAN 117, 103 (1957) (Ref.1) an approximate equation of the isothermal line of monomolecular adsorption with existing interactions adsorbate-adsorbate was given which describes the concave as well as the convex (and S-shaped) isothermal lines:

$$h = \theta' / K_1' (1 - \theta') (1 + K_n \theta'),$$

where h denotes the relative pressure of vapor, θ the degree of filling, K_1' , K_n the constants of equilibrium adsorbate-adsorbent and adsorbate-adsorbate. For the polymolecular adsorption (in Reference 2) equations were given which hold only in special cases: great K_1' with small K_n and small K_1' with great K_n . The authors here discuss a general case of

Card 1/4

Letter to the Editor

62-52-4-29/32

adsorption (polymolecular adsorption) for random K'_1 and K_n which taking into account that only the concentration of some uncoated complexes of the first layer enter the equilibrium equation (of the primary interaction adsorbate-adsorbent

$$K'_1 = \theta_{o1}/h(1 - \theta').$$

As with random

$$h\theta'_{o1}/\theta'_1 = \theta'_{o2}/\theta'_2 = \dots \theta'_o/\theta'$$

and with θ'_{o1} , θ'_{o2} and θ'_1, θ'_2 only the uncoated (and correspondingly to all single double and triple, and so on) horizontal complexes belong to the first layer, but θ'_n and θ' belong to all uncoated and to any other complexes of the first layer. Granted that the interactions adsorbate-adsorbate in the first layer do not depend on the fact whether the horizontal

Card 2/4

Letter to the Editor

62-58-4-29/52

complexes of this layer carry vertical complexes or not, according to reference 1 there is

$$\theta'_1 = \theta'_1 / 1 + K_n \theta'$$

the general degree of filling, however,

$$\theta = \theta'_1 / (1 - h)^2 = \theta' / (1 - h).$$

By introducing θ'_{o1} , θ'_1 , θ'_o and θ' into (2) the equation of the isothermal line of polymolecular vapor adsorption is obtained. Here the interaction adsorbate-adsorbate in the first layer is taken into account;

$$h = \frac{\theta(1 - h)^2}{K_1[1 - \theta(1 - h)][1 + K_n \theta(1 - h)]}.$$

At $K_n = 0$ this equation passes over into the equation BET (BET) just as well as the equation (Reference 1) into

Card 3/4

Letter to the Editor

62-58-4-29/32

Lanmuir's equation. There is 1 Soviet
reference.

ASSOCIATION: Institut fizicheskoy khimii Akademii nauk SSSR (Institute
of Physical-Chemistry, AS USSR)

SUBMITTED: February 15, 1958

AVAILABLE: Library of Congress

1. Polymolecular vapor absorption--Theory

Card 4/4

KISELEV, A.V.; POSHKUS, D.P.

Calculating the energy of adsorption of hydrocarbons on magnesium
oxide [with summary in English]. Zhur.fiz.khim. 32 no.12:2824-2834
D '58. (MIRA 12:2)

1. Moskovskiy gosudarstvennyy universitet imeni M.V. Lomonosova i
AN SSSR Institut fizicheskoy khimii, Moskva.
(Heat of adsorption) (Hydrocarbons) (Magnesia)

AUTHORS: Kiselev, A. V., Poshkus, D. P. SOV/20-120-4-40/67

TITLE: The Energy of the Coulomb Interaction Between the Hydroxyl Group of Silica Gel and the Benzene Molecule (Energiya kulonovskogo vzaimodeystviya gidroksil'noy gruppy silikagelya s molekuloy benzola)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol. 120, Nr 4, pp. 834 - 837 (USSR)

ABSTRACT: In the computations discussed in this paper the hydroxyl group is considered, to be a system of two point charges the position of which agrees approximately with the position of the oxygen atom and of the hydrogen atom. A formula is written down for the energy of the Coulomb (Kulon) interaction between the hydroxyl group and the benzene molecule; this energy was computed for various subcases in a vertical position of the hydroxyl group with regard to the direction of the benzene ring. The same energy in the case of a displacement of the hydroxyl group inside the benzene ring depends only in a low degree on their mutual position and amounts to about 4-6 kcal/mol. This interaction energy decreases considerably outside the benzene

Card 1/3

The Energy of the Coulomb Interaction Between the
Hydroxyl Group of Silica Gel and the Benzene Molecule

SOV/20-120-4-40/67

ring. According to the data obtained, the energy of the Coulomb interaction between the polar hydroxyl group and the non-polar benzene molecule is high and in the formation of the hydrogen binding in a given system it plays the decisive role among the other interactions. The benzene molecule adsorbed on the surface of a completely hydrated silica gel interacts essentially with one or with only a few hydroxyl groups; besides, such an adsorbed benzene molecule also interacts with the other atoms of the lattice of the silica gel. Therefore the removal of the hydroxyl groups from the surface of the silica gel is bound to decrease the heat of adsorption of the benzene molecules on the dehydrated surface of the silica gel. Besides, the energy of the interaction of the adsorbed benzene molecule must be increased with the volume phase. The computed energy of the Coulomb interaction (about 4-6 kcal/mol) agrees satisfactorily with the experimental values of the decrease of the adsorption heat of benzene in the dehydratization of the surface of silica gel. There are 1 figure and 20 references, 7 of which are Soviet.

Card 2/3

The Energy of the Coulomb Interaction Between the
Hydroxyl Group of Silica Gel and the Benzene Molecule

SOV/20-120-4-40/67

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im.M.V.Lomonosova
(Moscow State University imeni M.V.Lomonosov) Institut fiziches-
koy khimii Akademii nauk SSSR (Institute of Physical Chemistry
AS USSR)

PRESENTED: January 18, 1958, by A.N.Frumkin, Member, Academy of Sciences,
USSR

SUBMITTED: January 18, 1958

1. Hydroxyl radicals--Chemical reactions 2. Benzenes--Chemical re-
actions 3. Benzene molecules--Adsorption 4. Silicon dioxide--Chemical
properties 5. Chemical reactions--Energy

Card 3/3

5 (2), 5 (4)
AUTHORS:

Avgul' N. N., Kiselev, A. V.,
Lygina, I. A., Poshkus, D. P.

SOV/62-59-7-7/38

TITLE:

A Contribution to the Calculation of the Energy of the Adsorption
of Nonpolar Molecules on Graphite (K raschetu energii adsorbtsii
nepolyarnykh molekul na grafite)

PERIODICAL:

Izvestiya Akademii nauk SSSR. Otdeleniye khimicheskikh nauk,
1959, Nr 7; pp 1196-1206 (USSR)

ABSTRACT:

In this paper the details of a calculation of the adsorption
energies of simple and compound molecules carried out in a
previous paper are represented. The calculations were carried
out according to the formulas from paper reference 1 according to
which the adsorption energy is determined by the expressions

$$\Phi_i' = - C_{i1} \sum_j r_{ij}^{-6} - C_{i2} \sum_j r_{ij}^{-8} - C_{i3} \sum_j r_{ij}^{-10} + B_i' \sum_j e^{-r_{ij}/\rho}$$

$$\Phi_i'' = - C_{i1} \sum_j r_{ij}^{-6} - C_{i2} \sum_j r_{ij}^{-8} - C_{i3} \sum_j r_{ij}^{-10} + B_i'' \sum_j r_{ij}^{-12}$$

r_{ij} is the distance of the center of the i-th adsorption
molecule from the center of the j-th atom of the adsorbent.

Card 1/4

A Contribution to the Calculation of the Energy of the SOV/62-59-7-7/38
Adsorption of Nonpolar Molecules on Graphite

$C_{i1,2,3}$ are constants of the dispersion reaction. B' and B'' are the constant of the exponential function and the constant of the repulsion preceding the powers. g is an exponential constant of the repulsion. The calculation is carried out in two parts, the geometric one in which the distances of the adsorbed link i to all atoms j of the lattice of the adsorbent are calculated for different distances of the former from the surface. For this calculation only the lattice constants of the adsorbent have to be known. For the second part of the calculation of the forces the constants characterizing the reactions of both substances have to be determined. The calculation of the sums of r_{ij} was carried out for $n = 6, 8, 10$ and 12 for the different distances of the adsorbed link from the basis of the adsorbent equal to $2, 2.5, 3, 3.5$ a (a is the distance of the nearest atom). In table 1 the results of the calculation of the sums

$\sum_{ij}^{r_i} r_{ij}^{-n}$ are combined. The distances of the remaining graphite volume were determined from the integrals (3), (4), (5) (Table 2).

Card 2/4

A Contribution to the Calculation of the Energy of the Adsorption of Nonpolar Molecules on Graphite SOV/62-59-7-7/38

The sums $\sum_i^{r_i} r_{ij}^{-n} + \int_{(n)}$ are represented in tables 4 and 5 and the graphic representation in figure 2. The value $\sum e^{-r_{ij}/0.28}$ for g in table 6 was equated to 0.28 according to reference 6. For the sums of tables 4, 5 a simplified form with the constants p_n and q_n , the values of which are given in table 7, is introduced and the functions (1) and (2) are represented in the variable z

(6), (7). $\left(\sum_i^{r_i} r_{ij}^{-n} + \int_n = p_n z^{-q_n} \right)$. Next, the calculation of the reaction constant $C_{i1,2,3}$ is carried out. The values for different adsorptives are given in table 7 with the constants α and χ (polarizability, magnetic susceptibility) being necessary for the calculation of $C_{i1,2,3}$. ϕ' and ϕ'' were then determined by the aid of computed constants. The results for ϕ''

Card 3/4

A Contribution to the Calculation of the Energy of the Adsorption of Nonpolar Molecules on Graphite SOV/62-59-7-7/38

are given in tables 10 and 11. The calculation of the attraction- and repulsion constants was carried out from the balance energy of the adsorption of compound molecules on the basis of an additive scheme. There are 6 figures, 11 tables, and 8 references, 2 of which are Soviet.

ASSOCIATION: Institut fizicheskoy khimii Akademii nauk SSSR (Institute of Physical Chemistry of the Academy of Sciences, USSR)
Khimicheskiy fakul'tet Moskovskogo gosudarstvennogo universiteta im. M. V. Lomonosova (Chemical Department of the Moscow State University imeni M. V. Lomonosov)

SUBMITTED: November 16, 1957

Card 4/4

68700

S/069/60/022/01/005/025

D034/D003

56
5.4700
5.4400

AUTHORS:

Kiselev, A.V. and Poshkus, D.P.

TITLE:

The Heat and Entropy¹ of Adsorption of Benzene and N-
-hexane Vapors on Magnesium Hydroxide

PERIODICAL:

Kolloidnyy zhurnal, 1960, Vol XXII, Nr 1, pp 25-30
(USSR)

ABSTRACT:

The authors report on a study intended to determine the heat and entropy of adsorption of benzene and n-hexane vapors on magnesium hydroxide with weakly dehydrated surface. As adsorbed the authors selected a $Mg(OH)_{2-3}$ specimen moderately freed of water at a temperature of $20^{\circ}C$. The differential heat of adsorption Q_a was determined from the isotherms of adsorption on the given specimen. The isotherms were determined at the temperatures T_1 and T_2 (10 and $30^{\circ}C$) according to the formula

Card 1/4

Card 2/4

68700

S/069/60/022/01/005/025
D034/D003

The Heat and Entropy of Adsorption of Benzene and N-hexane Vapors on
Magnesium Hydroxide

be determined from isotherms as presented in a previous publication of the authors [Ref. 2]. The investigation has shown that the standard heat of adsorption of benzene vapors on the $\text{Mg}(\text{OH})_2$ specimen is greater than the standard heat of adsorption of n-hexane vapors (graph 1). A comparison of the standard heats of adsorption on various adsorbents (table and graph 4) makes evident that the heat of adsorption of benzene vapors on polar adsorbents is greater or close to the heat of adsorption of hexane, whereas on a non-polar adsorbent the heat of adsorption of benzene is considerably below the heat of adsorption of n-hexane. The increase in the ratio of the heats of adsorption of benzene and n-hexane vapors on polar adsorbents as compared to this ratio on a non-polar

Card 3/4

S/076/60/034/012/001/027
B020/B067

AUTHORS: Poshkus, D. P. and Kiselev, A. V.
TITLE: Energy of Dispersion Interaction of Benzene and n-Hexane
With the Surface of Magnesium Hydroxide
PERIODICAL: Zhurnal fizicheskoy khimii, 1960, Vol. 34, No. 12,
pp. 2640-2645

TEXT: In a preceding paper (Ref. 1), the authors assumed that the change in the ratio between the adsorption heats of benzene and n-hexane vapors on polar adsorbents is mainly due to the stronger electrostatic interaction of the benzene molecules as compared with the n-hexane molecules. Thus, an electric field is generated above the surface of the polar adsorbents, mainly above the lattice of $Mg(OH)_2$. Therefore, the authors studied the energy of interaction forces between benzene and n-hexane molecules and the $Mg(OH)_2$ surface. $Mg(OH)_2$ has a multi-layer rhombohedral ionic crystal lattice of the type CsI_2 . Each lattice layer consists of an

✓

Card 1/3

Energy of Dispersion Interaction of Benzene and n-Hexane With the Surface of Magnesium Hydroxide S/076/60/034/012/001/027 B020/E067

Mg ion plane which lies between two hydroxyl ion planes which, in turn, consist of oxygen and hydrogen ion planes. The Φ_{1D} potential of dispersion interaction of the member i of the chain of hydrocarbon molecules (CH_3 , CH_2 and CH_{arom}) with the centers of force j of the $\text{Mg}(\text{OH})_2$ lattice (H, O, and Mg) was calculated for three layers above the basal plane of $\text{Mg}(\text{OH})_2$, i.e., 1) above the hydroxyl ion of the first (external) plane (layer A), 2) above the center of the hydroxyl-ion triangle of the first plane, below which lies the magnesium ion of the second plane (layer B), or 3) the hydroxyl ion of the third plane (layer C). When expanding the dispersion potential in a series, only the first two terms are considered which represent the potential of dipole - dipole and the dipole - quadrupole interactions. The others influence the entire potential only slightly. The constants of dispersion interaction were calculated. The sums of the exponential functions $\sum r_{ij}^{-n}$, where $n = 6$ and 8 , and $j = \text{H, O, or Mg}^{2+}$, were calculated by assuming that $a = 3.11 \text{ \AA}$, $c = 4.73 \text{ \AA}$, $u = 0.22$, and that the distance

Card 2/3

Energy of Dispersion Interaction of Benzene and n-Hexane With the Surface of Magnesium Hydroxide S/076/60/034/012/001/027 B020/B067

between the hydrogen and oxygen atom centers in the hydroxyl ion is $l_{OH^-} = 0.97$ Å. Summation was performed up to $r_{ij} = 10$ Å. Hence, the great increase in the ratio between the adsorption heats of benzene and n-hexane on $Mg(OH)_2$ as compared to those of the vapors of these

compounds on graphite is not caused by the dispersion interaction of benzene and n-hexane molecules with the basal plane of $Mg(OH)_2$. There are 3 figures, 5 tables, and 15 references: 6 Soviet, 2 US, 3 British, and 4 German.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M. V. Lomonosova (Moscow State University imeni M. V. Lomonosov). Akademiya nauk SSSR, Institut fizicheskoy khimii (Academy of Sciences USSR, Institute of Physical Chemistry) ✓

SUBMITTED: October 25, 1958

Card 3/3

KISELEV, A.V.; POSHKUS, D.P. [Poskus, D.]

Molecular-statistical calculation of the heat capacity of argon and benzene adsorbed on graphite. Zhur.fiz.khim. 37 no.7:1504-1509 J1
'63. (MIRA 17:2)

1. Institut fizicheskoy khimii AN SSSR i Institut khimii i khimicheskoy tekhnologii AN Litovskoy SSR.

POSHKUS, D.P. (Moscow); KISELEV, A.V. (Moscow)

Electrostatic field above the basal plane of magnesium hydroxide,
and its interaction with benzene and n-hexane molecules. Zhur.
fiz. khim. 34 no.12:2647-2653 D '60. (MIRA 14:1)

1. Institut fizicheskoy khimii AN SSSR i Moskovskiy gosudarstvennyy
universitet imeni M.V. Lomonosova.
(Magnesium hydroxide) (Benzene) (Hexane)

KISELEV, A.V.; POSHKUS, D.P. [Poskus, D.]

Statistical thermodynamic calculation of adsorption
equilibrium for benzene on graphite. Dokl. AN SSSR
139 no.5:1145-1148 Apr '61. (MIRA 14:8)

1. Institut khimii i khimicheskoy tekhnologii AN Litovskoy SSR
i Institut fizicheskoy khimii AN SSSR. Predstavleno akademikom
A.N. Frumkinym.

(Benzene) (Adsorption)

POSHKUS, D.P.; KISELEV, A.V.

Contribution to the statistical thermodynamic calculation of
adsorption equilibrium in the system argon - graphite. Zhur.fiz.
khim. 36 no.8:1735-1742 Ag '62. (MIRA 15:8)

1. Institut khimii i khimicheskoy tekhnologii AN Litovskoy SSR i
Gruppa khimii poverkhnosti Instituta fizicheskoy khimii AN SSSR.
(Argon) (Graphite) (Adsorption)

VALENTELIS, L. Yu.; REKLITE, V.V. [Raklyte, V.]; POSHKUS, D.P. [Poskus, D.]
MATULIS, Yu.Yu. [Matulis, J.]

Correlation between texture and hydrogen absorption by nickel
electrodeposits as dependent on the conditions of electrolysis.
Trudy AN Lit. SSR. Ser. B. no. 4:7-14 '65 (MIRA 19:2)

1. Institut khimii i khimicheskoy tekhnologii AN Litovskoy SSR.
Submitted July 29, 1965.

POSHKUS, D.P. [Poskus, D.]; KAZLAUSKAS, R.M.

Adsorption on silica gel of methylene blue from aqueous solutions.
Koll. zhur. 27 no.5:738-744 S-O '65. (MIRA 18:10)

1. Institut khimii i khimicheskoy tekhnologii AN Litovskoy SSR.

KISELEV, A.V.; POSHKUS, D.P. [Poskus, D.]; AFREYMOVICH, A.Ya.

Theoretical calculation of the potential function and thermodynamic properties of symmetric diatomic molecules adsorbed on graphite nitrogen adsorption. Zhur. fiz. khim. 39 no.5:1190-1197 My '66.
(MIRA 18.8)

1. Institut fizicheskoy khimii AN SSSR i Institut khimii i khimicheskoy tekhnologii AN Litovskoy SSR.

KITSELEV, A.V.; POSHKUS, D.P. [Poskus, D.P.]; AFREYMOVICH, A.Ya.

Molecular statistical calculation of the thermodynamic properties
of inert gases adsorbed on graphite. Izv. Akad. Nauk, 1964, no. 11:
1514-1522. Je '64. (MIRA 18:3)

1. Institut fizicheskoy khimii AN SSSR i Institut khimii i khimicheskoy tekhnologii Akademii nauk Litovskoy SSR.

KISELEV, A.V.; POSHKUS, D.P. [Poskus, D.]

Theoretical calculation of the potential function and the thermodynamic properties of symmetrical diatomic molecules adsorbed on graphite. Adsorption of H_2 and D_2 . Zhur. fiz. khim. 39 no.2:398-402 F '65. (MIRA 18:4)

1. Institut fizicheskoy khimii AN SSSR i Institut khimii i khimicheskoy tekhnologii AN Litovskoy SSR.

KISELEV, A.V.; POSHKUS, D.P. [Poskus, D.P.] (Moscow)

Statistical calculation of the total energy and entropy of
argon adsorbed on graphite. Zhur. fiz. khim. 37 no.4:770-777
Ap '63. (MIRA 17:7)

1. Akademiya nauk SSSR, Institut fizicheskoy khimii i Institut
khimii i khimicheskoy tekhnologii AN Litovskoy SSR.

KISELEV, A. V.; POSHKUS, D. P.

Statistical calculation of the total energy and entropy of
benzene adsorbed on graphite. Zhur. fiz. khim. 37 no. 3:602-
614, Mar '63. (MIRA 17:5)

1. Institut fizicheskoy khimii AN SSSR i Institut khimii i khim-
icheskoy tekhnologii AN Litovskoy SSSR.

L 1691/-63
WM/JD/JW/K

EPR/EPF(c)/EWP(q)/EWT(m)/BDS AFFTC/ASD Ps-4/Pr-4 WE/
S/076/63/037/004/005/029

AUTHOR: Kiselev, A. V., Poshkus, D. P.

TITLE: Statistical calculation of the total energy and entropy of argon
adsorbed on graphite 72
10 27

PERIODICAL: Zhurnal fizicheskoy khimii, V. 37, No. 4, 1963, 770-776

TEXT: A statistical calculation was made of the total energy and entropy of argon which was adsorbed on a graphite surface. The procedure was based on the calculation of the potential energy of argon molecules at the graphite surface using the approximate theory of intermolecular interactions. The effect of various approximations of this function on the values computed for the thermodynamic functions of adsorbed argon was investigated. Expressions were obtained for changes in the thermodynamic functions of argon upon adsorption on graphite in a classical approximation and with an approximate reckoning of the quantum mechanical corrections. The calculated theoretical values for the change in total energy and entropy of argon adsorbed on graphite were in good accord with values which were obtained experimentally using graphitized carbon black. This is similar to the results which were obtained by the authors in a previous work for the case of the adsorption of benzene on graphite. There are 3 tables and a graph. The most

Card 1/2

L 16914-63

S/076/63/037/004/005/029

Statistical calculation of the total energy ...

2

important English-language reference reads as follows: S. Ross, W. J. Winkler, J. Coll. Sci., 10, 319, 1955.

ASSOCIATION: Akademiya nauk SSSR, Institut fizicheskoy khimii, Institut khimii i khimicheskoy tekhnologii AN LitSSR (Academy of Sciences of the USSR, Institute of Physical Chemistry, Institute of Chemistry and Chemical Technology of the Academy of Sciences Lithuanian SSR)

SUBMITTED: February 22, 1962

Card 2/2

I: 17728-63 EPR/EPF(c)/EWT(1)/EPF(n)-2/EWP(q)/EWT(m)/RDS AFFTC/
AND/SSD P=4/PL=4 P=4/PL=4 WH/NW/JN/JD/K

ACCESSION NR: AP3004060

S/0076/63/037/007/1504/1509

AUTHORS: Kiselev, A. V.; Poskus, D. P.

TITLE: Molecular-statistical calculation of heat capacity of argon and benzene
adsorbed on graphite

SOURCE: Zhurnal fizicheskoy khimii, v. 37, no. 7, 1963, 1504-1509

TOPIC TAGS: benzene, argon, graphite, statistical heat capacity calculation,
molecular heat capacity calculation

ABSTRACT: This is a continuation of a series of analyses which the authors carried out with respect to calculation of the full energy, entropy and chemical potential of argon and benzene which were adsorbed on the basal face of graphite at low surface coverage, proceeding from the interaction of the molecules of the adsorbate with the atoms of the adsorbent. In this work, authors carry out an analogous molecular-statistical calculation of the heat capacity of argon and benzene on graphite. Treatment is mathematical and results are tabulated. The molecular-statistical calculations were made for the differential molar heat capacities of isolated argon and benzene molecules adsorbed on a uniform basal face of graphite. Authors conclude that the calculated values for the heat capacities of monoatomic

Card 1/2

L 17728-63

ACCESSION NR: AP3004060

2

molecules of argon do not overlap for models of localized and non-localized adsorption for various force constants of hindered molecular motion along the surface. This fact may be used for an unequivocal determination of the state of the adsorbed monoatomic molecules from experimental data obtained at low surface coverage. Orig. art. has: 13 tables and 16 formulas.

ASSOCIATION: Akademiya nauk SSSR, Institut fizicheskoy khimii (Academy of sciences, SSSR, Institute of physical chemistry); Akademiya nauk LitSSR, Institut khimii i khimicheskoy tekhnologii (Academy of sciences, Lithuanian SSR, Institute of chemistry and chemical engineering)

SUBMITTED: 20 June 62

DATE ACQ: 15 Aug 63

ENCL: 00

SUB CODE: PH, CH

NO REF SCV: 011

OTHER: 005

Cord 2/2

S/076/63/037/003/010/020
B101/B215

AUTHORS: Kiselev, A. V., Poshkus, D. P. (Moscow)

TITLE: Statistical calculation of the total energy and entropy of
benzene adsorbed by graphite

PERIODICAL: Zhurnal fizicheskoy khimii, v. 37, no. 3, 1963, 608-614

TEXT: The statistical calculation of the total energy and entropy of
adsorbed benzene at a low occupation θ of the surface was based on a
calculation of the potential energy of the benzene molecule adsorbed by
a graphite surface by using the approximate molecular interaction theory.

The mathematical apparatus of these calculations is based on a
publication by the author in Trans. Faraday Soc., 59, 428, 1963, and on a
paper by J. W. Dreenan, T. L. Hill (J. Chem. Phys. 17, 775, 1949).
Results:

Card: 1/3

Statistical calculation of the total ...

S/076/63/037/003/010/020
B101/B215

$$U_{a \text{ кл.}} = \Phi_{00} + RT \left[\frac{7}{2} + \frac{\int_0^{d/z_{00}} \exp(-\Delta\Phi_0/kT) (\Delta\Phi_0/kT) (kT/h\nu_z) f(\beta) d\beta}{\int_0^{d/z_{00}} \exp(-\Delta\Phi_0/kT) (kT/h\nu_z) f(\beta) d\beta} \right] \quad (12)$$

$$S_{a \text{ кл.}} = R \left[\frac{7}{2} + \ln \frac{8\pi^3}{\sigma} \frac{2\pi mkT}{h^3} \frac{2\pi AkT}{h^3} \left(\frac{2\pi CkT}{h^3} \right)^{1/2} \frac{z_{00} \omega_m}{d} \times \right. \\ \left. \times \int_0^{d/z_{00}} \exp\left(-\frac{\Delta\Phi_0}{kT}\right) \frac{kT}{h\nu_z} f(\beta) d\beta + \right. \\ \left. + \frac{\int_0^{d/z_{00}} \exp(-\Delta\Phi_0/kT) (\Delta\Phi_0/kT) (kT/h\nu_z) f(\beta) d\beta}{\int_0^{d/z_{00}} \exp(-\Delta\Phi_0/kT) (kT/h\nu_z) f(\beta) d\beta} \right] - R \ln \theta. \quad (13)$$

where Φ_{00} is the adsorption heat of benzene at 0°K , $\theta \rightarrow 0$; the indices кл. stand for classical. For adsorbed benzene at 293°K , the calculation yields Card 2/3

Statistical calculation of the total ...

S/076/63/037/003/010/020
B101/B215

a total energy $U_{a\text{ cl}} = -6.93$ kcal/mole and entropy $S_{a\text{ cl}} = 39.01 - 4.57 \log \theta$ entropy units. After subtracting the data for gaseous benzene at 293°K : $U_g = 1.75$ kcal/mole, $S_g = 59.02$ e. u., the following values are obtained $U_{a\text{ cl}} = -8.68$ kcal/mole, $S_{a\text{ cl}} = -20.01 - 4.57 \log \theta$ e. u. After quantum mechanical correction, $\Delta U_a = -8.68$ kcal/mole, $\Delta S_a = -19.9 - 4.57 \log \theta$ e. u. are obtained under consideration of the approximations by K. S. Pitzer and W. D. Gwinn (J. Chem. Phys., 10, 428, 1942). The data are in good agreement with the experimental data obtained for benzene adsorbed by graphited carbon black. There are 1 figure and 3 tables.

ASSOCIATION: Institut fizicheskoy khimii AN SSSR (Institute of Physical Chemistry AS USSR); Institut khimii i khimicheskoy tekhnologii AN LitSSR (Institute of Chemistry and Chemical Technology of AS LitSSR)

SUBMITTED: February 22, 1962

Card 3/3

POSHKUS, N.B.

Effect of some hormonal preparations on blood coagulation and pro-
thrombin time in rabbits. Trudy Ukr. nauch.-issl. inst. eksper. endok.
19:397-403 '64. (MIRA 18:7)

1. Iz otdela patofiziologii Ukrainskogo instituta eksperimental'noy
endokrinologii i kafedry fizioterapii Ukrainskogo instituta usovershen-
stvovaniya vrachey.

POSHKUS, N.B.

Effect of hexonium on blood coagulation and the prothrombin
time in rabbits. Trudy Ukr.nauch.-issl.inst.eksper.endok.
18:369-374 '61. (MIRA 16:1)
(PROTHROMBIN) (BLOOD--COAGULATION) (HEXONIUM)

POSHOGINA, M.

Characteristics of the liver of the Baltic codfish. Rybnoye Khoziaystvo
26, 41-3 '50. (MLBA 3:1)
(CA 47 no.18:9566 '53)

POSEPUR, A.A., NECHIPORCHUK, I.D., doktor sel'khoz. nauk

Making use of the qualitative differences of hop tissues in hop production. Agrobiologiya no.5:761-763 3-0 '60. (MIRA 13:10)

1. Sel'skokhozyaystvennyy institut, L'vov.
(Hops)

POSHUMENSKIY, Semen Vladimirovich; MEZENTSEV, S., red.; TSIVUNIN, I.,
tekhn. red.

[Energetic start] Energichnyi start. Syktyvkar, Komi knizhnoe
izd-vo, 1960. 57 p. (MIRA 14:9)
(Komi A.S.S.R.—Socialist competition)

AUTHORS: Posik, L.N., Tenenbaum, I.M.

89-7-5/32

TITLE: The Application of a Special Apparatus for Express Analysis of Mined Ores by Means of γ -Radiation (Primeneniye spetsial'noy apparatury dlya ekspress-analizov dobytykh rud po γ -izlucheniyu)

PERIODICAL: Atomnaya Energiya, 1957, Vol. 3, Nr 7, pp. 28-35 (USSR)

ABSTRACT: For the operative computation of the amount and the quality of the mined ores and of the production in various containers (wagons, trucks, chests) and also for the separation of the rock enriched with uranium, the filling material, etc., the PKC device (station-control radiometer) is widely used. The present paper describes these devices as well as the mining-biological conditions for their use. Also these problems are mentioned which can be solved by means of this device. From the point of view of the rational application of the devices PKC 1 and PKC 2 for the operative calculation of mining for the estimation of production, and for measures to be taken against loss of metal as well as against the impoverishment of the ore, all ore mines can be subdivided into two groups: 1.) Such with a summary working of the ore, 2.) Such with selective working of the ore. In mines with summary working the entire

Card 1/2

The Application of a Special Apparatus for Express
Analysis of Mined Ores by Means of γ -Radiation

89-7-5/32

operative calculation of the mined ore and of the ore destined for commerce is based upon the γ -express analyses of the entire mine quantity in wagons and trucks by means of the PKC-2 devices. In the case of selective working estimation of the mined ore is carried out with the PKC-1 devices. This calculation serves as a basis for the determination of the mined quantities in the case of hydrothermal vein-like deposits with rich ore bodies of small thickness. Also the express analysis of rich commercial ores is carried out by means of devices of the PKC-1 type. By means of the devices PKC-2 it is possible not only to determine the activity of ores and their nature in an operative manner in the individual containers, but also to automatize the tedious work of sorting-in and discharging from trucks. There are 8 figures, 1 table, and 3 references, 3 of which are Slavic.

SUBMITTED: October 9, 1956

AVAILABLE: Library of Congress

Card 2/2

1. Radiometers - Applications
2. Ores (Radioactive) - Analysis - Equipment
3. Mining engineering - USSR

AUTHOR: Posik, L. N.

89-4-5-12/26

TITLE: The Radiation Field of a Rectangular Parallelepiped, **Taking Self-Absorption Into Account**
(Pole izlucheniya pryamougol'nogo parallelepipeda s uchetom samopogloshcheniya)

PERIODICAL: Atomnaya Energiya, 1958, Vol. 4, Nr 5,
pp. 470 - 471 (USSR)

ABSTRACT: For the construction of an apparatus and of a method for the express analysis of rocks in a reservoir the above mentioned field had to be determined. Under the presumption that the spectral composition of the γ - emitter is not taken into consideration, the γ -dose in point A (at the same time also origin of coordinates) can be computed from the equation:

$$P = K_{qv} \int_0^m dx \int_0^n dy \int_0^1 \frac{e^{-\mu \sqrt{x^2 + y^2 + z^2}}}{x^2 + y^2 + z^2} dz$$

Card 1/2

89-4-5-12/26

The Radiation Field of a Rectangular Parallelepiped, Taking Self-Absorption Into Account

K = ionization constant of the active substance

q_v = volume concentration of the substance

m, n, l = the lateral lengths of the parallelepiped

μ = the γ -absorption coefficient.

It is shown that it is possible to represent the triple integral as a sum combination of the simple integrals. The numerical result for 2 cases with 6 different μ values is given. There are 1 figure, 1 table and 10 references, 7 of which are Soviet.

SUBMITTED: January 15, 1958

AVAILABLE: Library of Congress

1. Radiation--Mathematical analysis 2. Recks--Radioactivity--Measurement

Card 2/2

POSIK, Lev Notovich; KOSHKELEV, Ivan Vasil'yevich; BOVIN, Vladimir Pavlovich; SAGURO, M.A., red.; MAZEL', Ye.I., tekhn.red.

[Rapid radiometric determination of mined ores; brief guide]
Radiometricheskii ekspress-analiz dobytykh rud; kratkoe rukovodstvo. Moskva, Izd-vo Glav.upr. po ispol'zovaniyu atomnoi energ. pri Sovete Ministrov SSSR, 1960. 75 p.

(MIRA 13:12)

(Uranium ores) (Radioactivity--Measurement)
(Ores--Sampling and estimation)

SURAZHSKIY, Daniil Yakovlevich. Prinimeli uchastiye: PUKHAL'SKIY, L.Ch.;
POSIK, L.N.; SHASHKIN, V.L.. SMIRNOV, V.I., red.; ALYAB'YEV, A.F.,
red.; POPOVA, S.M., tekhn.red.

[Methods of prospecting and exploration of uranium deposits]
Metody poiskov i razvedki mestorozhdenii urana. Pod red. V.I.
Smirnova. Moskva, Izd-vo glav.upr.po ispol'zovaniyu atomnoi
energii pri Sovets Ministrov SSSR, 1960. 240 p.

(MIRA 13:7)

1. Chlen-korrespondent AN SSSR (for Smirnov).
(Prospecting) (Uranium ores)

POSIK, L.N.; BIBICHENKO, S.I.; GRODKO, R.A.

[Radiometric analysis of ores on conveyers] Radiometricheskii analiz rud na transporterakh. Moskva, Glav. upr. po ispol'zovaniyu atomnoi energii, 1960. 18 p.
(MIRA 17:1)
(Ores—Radioactive properties) (Radiometry)

KADLUBOWSKI, R.; POSIŁA, L.

On the third fraction of coagulating blood. Acta physiol. polon 3
Suppl. 3: 167-168 1952. (CLML 24:1)

1. Of the Institute of General Pathology (Head--Prof. F. Vermet, M.D.)
of Lodz Medical Academy.

KADLUBOWSKI, Roscislaw; POSILA, Leszek, Lodz.

Identity of Danilin and Biernacki reactions. Przegl.lek., Krakow,
11 no.9:274-277 1955.

1. Z Zakladu Patologii Ogolnej A M w Lodzi. Kierownik: prof. dr.
F. Venulet.

(BLOOD SEDIMENTATION

Biernacki, & Danilin reactions, comparison)

POSINKOVIC, B.

Dyschondroplasia--Ollier's disease. Acta chir. Iugo. 8 no.3:
253-261 '61.

1. Bolnica za tuberkulozu kostiju i zglobova, Biograd n/m (Ravnatelj
dr B. Metz).
(DYSCHONDROPLASIA)

POSINKOVIC, Bozidar, dr.

Bone tuberculosis in Dalmatia. Lijecn. vjesn. 87 no.6:611-617
Je '65.

1. Iz Ortopedske bolnice u Biogradu n/m.

POSINKOVIC, Marko, dr

Integration movement in construction engineering.
Gradevinar 16 no.3:113-114 Mr '64.

DUANCIC, Vjekoslav; POSINOVEC, Jasminka

On the nature of the parotid gland in adults. Rad. med. fak. Zagreb.
10:215-222 '62.

(PAROTID GLAND)

S

YUGOSLAVIA

Vjekoslav DUANCIC and Jasminka POSINOVEC, Department of Histology and Embryology of Medical Faculty (Zavod za histologiju i embriologiju) University of Zagreb.

"Type of Alveoli of Parotid Gland in Adults."

Zagreb, Radovi Medicinskog Fakulteta u Zagrebu, Vol 10, No 3, 1962; pp 215-222.

Abstract [French summary modified]: Studies on 25 male and 27 female cadavers (9120 slides) revealed that the universally found statements contending that the parotid gland has only serous alveoli is in error: mucous alveoli were found in 9 of the male and 1 female specimen. Two tables, 48 references mostly histologic and other textbooks - 44 Western, 2 Yugoslav, Czech, Soviet.

1/1

POSINOVEC, Jasminka

A rare discovery of sebaceous gland in the parotid gland. Rad. med.
fak. Zagreb. 10 no.2:163-173 '62.

(PAROTID GLAND)

(SEBACEOUS GLAND)

POSIPISIL, F.; PRAT, S.

Humic acids with C^{14} . In English. p. 71.

BIOLOGIA PLANTARUM. (Ceskoslovenska akademie ved. Biologicky ustav)
Praha, Czechoslovakia, Vol. 1, no. 1, 1959.

Monthly List of East European Accessions (EEA^T), LC, Vol. 8, no. 11, Nov. 1959
Uncl.

1ST AND 2ND COLUMNS																										3RD AND 4TH COLUMNS																									
1ST AND 2ND COLUMNS													3RD AND 4TH COLUMNS																																						
<p>CA</p> <p>9</p> <p>MOVEMENT AND DISTRIBUTION OF BURDEN IN THE BLAST FURNACE. M. B. POIN. <i>Domes</i> 1934, No. 1, 19-27. A theoretical discussion of the effect of distribution of materials in the blast furnace on its performance. In large furnaces, rate of descent of charge increases from the center outward. Concentration of ore in the charge increases in the same direction. A uniform distribution of the ore through the cross-section of the furnace will not increase the capacity of the furnace or decrease coke consumption. Since the gas passing through the charge increases in temp. from the center outward, the operation of the furnace should be so controlled that the outer ring of charge should descend at a greater rate than the inner rings. This can be accomplished by sorting the ore and coke as they are thrown from the hopper in such a way that lump size should increase from the center outward, and be smaller again at the wall. The sorting app. developed by Sorokin is well adapted for this purpose. S. L. M.</p>																																																			
<p>ASTM-SL-4 METALLURGICAL LITERATURE CLASSIFICATION</p>																																																			

POSIN, N. V.

"Telephasometer" (Telefazometr) from the book Telemechanization in the National Economy, pp. 310-314, Iz. AN SSSR, Moscow, 1956

(Given at meeting held in Moscow 29 Nov to 4 Dec 54 by Inst. of Automatics and Telemechanics)

POSH, S. S., YURGEV, V. Y., SKURICHINA, G. M., BILICH, I. N.

"Absorption of organic molecules in cellulose," a paper presented at the 9th Congress on the Chemistry and Physics of High Polymers, 28 Jan-2 Feb 57, Moscow, Textile Research Inst.

B-3,084,395

DUANCIC, Vj; POSINOVEC, J.

The atrial myocardium and the artrioventricular valves. Med.
arh. 15 no.3:13-24 My-Je '61.

1. Zavod za histologiju i embriologiju Medicinskog fakulteta u
Zagrebu (Predstojnik: prof. dr Vj. Duancic).
(HEART VALVES anat & histol)

POSINOVEC, Jasminka

Tissular mastocytes and some new concepts on them. Biol glas
15 no.2:127-132 '62.

1. Zavod za histologiju i embriologiju Medicinskog fakulteta
u Zagrebu.

STANESCU, Panait; MERCEA, Florian, ing., correspondent; POSIRCARU, Alexandru; AVACUM, Mihail

Technological tests will start 20 days earlier. Constr
Buc 16 no. 749:4 16 May '64.

1. Chairman of the Trade-Union, Committee of the Construction Site of Straw Paper and Cellulose concern Calarasi (for Stanescu).
2. The Galati Branch of the Voluntary Editorial Office of "Constructorul" (for Posircaru, Avacum).

DUANCIC, Vjekoslav; POSINOVEC, Jasminka

Contribution to the study of histological structure of auriculo-ventricular fibrous rings. Radovi med. fak., Zagrebu 7 no.1:57-62 '59.

(MYOCARDIUM anat. & histol.)

POSIRCARU, A., instalator; AVACUM, M., normator

Successes in production and cultural activity. Constr Buc
16 no.736:4 15 F'64.

POSIT, Z.

Thermoelectric phenomena and their applications. P 754

SLABOPROUDY OBZOR (Ministerstvo vsobenibo strojirenstvi, Ministerstvo spoju
a Ceskoslovenska vedecko-technicka spolecnost, sekce elektrotechnika) Praha,
Czechoslovakia, Vol. 20, no. 12 Dec. 1959

Monthly List of East European Accessions (EEAI). LC. Vol. 9, no. 2,
Feb. 1960

Uncl.

POSITAKO, I.

Method of joining high pressure pipes with cones. Rech. transp.
21 no. 4:30 Ap '62. (MIRA 15:4)

1. Pomoshchnik mekhanika zemsnyaya "DN-19" Dneprovskogo
tekhnicheskogo uchastka.
(Marine pipe fitting)

POSIVAL, V.

Sobotka, M.; Posival, V. Efficiency tests of circuit breakers. p. 139.
ELEKTROTECHNIK. Praha. Vol. 10, no. 5, May 1955.

SO: Monthly List of the East European Accession, (EEAL), LC. Vol. 4,
no. 10, Oct. 1955. Uncl.

POSIVAL, V.

Sobotka, M. Efficiency tests of circuit breakers. p. 139.
ELEKTROTECHNIK, Praha, Vol. 10, no. 5, May 1955.

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 4, no. 10, Oct. 1955,
Uncl.

POSKACHEY, A.A. (Moskva); SVET, D.Ya. (Moskva)

Investigating the radiating properties of aluminum alloys in the
near infrared region of the spectrum. Izv. AN SSSR. Otd. tekhn. nauk.
Met. i topl. no. 3:86-91 My-Je '60. (MIRA 13:6)
(Aluminum alloys--Thermal properties) (Spectrum, Infrared)

POSKACHEY, A.A.

Measuring temperatures under 800°C by radiation. ISvet. met. 36
no.9:76-79 S '63. (MIRA 16:10)

POSKACHEY, A.A.

Results of the study of germanium photodiodes. Trudy VZET
no.18:32-41 '61. (MIRA 17:1)

POSPACHENY, A.A. (Moskva); URAPOV, M.M. (Moskva)

Color temperature of aluminum and magnesium alloys in the
300-500° C range. Izv. AN SSSR Met. i gor. delo no.2:186-187
Mn-Ap'64 (MIRA 17:8)

L 24406-66 ENT(d)/EMP(h)/EMP(1)

ACC NR: AP6006370

SOURCE CODE: UR/0413/66/000/002/0106/0106

AUTHOR: Poskachev, A. A.

ORG: none

TITLE: Automatic multichannel spectral ratio pyrometer. Class 42, No. 178143

SOURCE: ¹⁴ Izobreteniya, promyshlennyye obraztsy, tovarnyye znaki, no. 2, 1966, 106

TOPIC TAGS: pyrometer, pyrometry, *optic pyrometer*

ABSTRACT: This Author Certificate presents an automatic multichannel spectral ratio pyrometer which contains detectors, shift and amplifier-converter devices, and a marker device (see Fig. 1). For simultaneous rapid-response recording of the temperatures of several objects on one photographic film, the pyrometer has a stroboscopic device in the form of a pulsed point light source and a mirror rotated synchronously by a motor. The light source is connected to the output of the amplifier-converter unit.

Card 1/2

UDC: 536.521.3

L 24406-66

ACC NR: AP6006370




Fig. 1. 1 - pulsed point light source; 2 - amplifier-converter unit; 3 - mirror; 4 - motor; 5 - pulling mechanism with photographic film.

Orig. art. has: 1 diagram.

SUB CODE: 20,14/ SUM DATE: 27Jul62

Card 2/2dda

L 24407-66

ACC NR: AP6006371

SOURCE CODE: UR/0413/66/000/002/0106/0107

AUTHOR: Poskachev, A. A.

ORG: none

TITLE: Spectral ratio pyrometer.¹⁰ Class 42, No. 178144

SOURCE: Izobreteniya, promyshlennyye obraztsy, tovarnyye znaki, no. 2, 1966, 106-107

TOPIC TAGS: pyrometer, pyrometry, *optic pyrometer*

ABSTRACT: This Author Certificate presents a spectral ratio pyrometer containing an optical system, two light filters, a shutter, frame windows, and a radiation receiver. To measure the ratio of radiation intensities, the pyrometer has two phase shift devices connected to the radiation receiver and a phase detector connected to the phase shift devices (see Fig. 1). A slave system connected to the output of the phase detector is mechanically coupled to one of the phase shift devices.

Card 1/2

UDC: 536.521.3

L 24407-66

ACC NR: AP6006371




Fig. 1. 1 and 2 - phase shift
devices; 3 - radiation receiver;
4 - phase detector.

Orig. art. has: 1 diagram.

SUB CODE: 20/4/

SUBM DATE: 27Jul62

Card 2/2 dda

L 24408-66

ACC NR: AP6006372

AUTHOR: Poskachev, A. A.

ORG: none

TITLE: Digital spectral ratio pyrometer. Class 42, No. 178145

SOURCE: Izobreteniya, promyshlennyye obraztsy, tovarnyye znaki, no. 2, 1966, 107

TOPIC TAGS: pyrometer, digital system, *electronic circuit, circuit design*

ABSTRACT: This Author Certificate presents a digital spectral ratio pyrometer containing a detector, a regulated phase shift device, and an amplifier-converter circuit. To decrease the time lag and to simplify the design, the pyrometer has a coincidence circuit with its input connected to an oscillator and (through the amplifier-converter device) to the detector (see Fig. 1). An electronic counter for converting the intensity ratio of two fluxes into digital code is connected to the output of the coincidence circuit.

UDK: 536.521.3

Card 1/2

L 24408-66
ACC NR: AP6006372




Fig. 1. 1 - Coincidence circuit; 2 - detector; 3 - amplifier-converter device; 4 - oscillator; 5 - electronic counter; 6 - phase shift device.

Orig. art. has 1 diagram.

SUB CODE: 20,14,01/SUBM DATE: 27Jul62

Card 2/2dda